



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### Drought-induced mortality in Scots pine

**Citation for published version:**

MacAllister, S, Mencuccini, M, Sommer, U, Engel, J, Hudson, A, Salmon, Y & Dexter, KG 2019, 'Drought-induced mortality in Scots pine: Opening the metabolic black box', *Tree physiology*, vol. 39, no. 8, tpz049, pp. 1358-1370. <https://doi.org/10.1093/treephys/tpz049>

**Digital Object Identifier (DOI):**

[10.1093/treephys/tpz049](https://doi.org/10.1093/treephys/tpz049)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Peer reviewed version

**Published In:**

Tree physiology

**Publisher Rights Statement:**

This is a pre-copyedited, author-produced version of an article accepted for publication in *Tree physiology* following peer review. The version of record Sarah MacAllister, Maurizio Mencuccini, Ulf Sommer, Jasper Engel, Andrew Hudson, Yann Salmon, Kyle G Dexter, Drought-induced mortality in Scots pine: opening the metabolic black box, *Tree Physiology*, , tpz049, <https://doi.org/10.1093/treephys/tpz049> is available online at: <https://academic.oup.com/treephys/advance-article/doi/10.1093/treephys/tpz049/5481586>

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



**Drought-induced mortality in Scots pine: opening the metabolic black box**

Sarah MacAllister<sup>1</sup>,  
Maurizio Mencuccini<sup>2,3</sup>,  
Ulf Sommer<sup>4</sup>,  
Jasper Engel<sup>4</sup>,  
Andrew Hudson<sup>5</sup>,  
Yann Salmon<sup>6</sup>,  
Kyle G. Dexter<sup>1,7</sup>

<sup>1</sup>School of GeoSciences, University of Edinburgh, UK

<sup>2</sup>CREAF, Barcelona, Spain

<sup>3</sup>ICREA, Barcelona, Spain

<sup>4</sup>NBAF-Birmingham, School of Biosciences, University of Birmingham, UK

<sup>5</sup>School of Biological Sciences, University of Edinburgh, UK

<sup>6</sup>University of Helsinki, Finland

<sup>7</sup>Royal Botanic Garden Edinburgh, UK

Corresponding Author: Sarah MacAllister

S.l.wilkinson2@gmail.com

Word counts for abstract (200) and main text (6463)

Number of tables (3)

Number of figures (3)

Number of references (69)

## ABSTRACT

- Forests are sensitive to droughts, which increase the mortality rate of tree species. Various processes have been proposed to underlie drought-induced tree mortality, including hydraulic failure, carbon starvation, and increased susceptibility to natural enemies. To give insights into these processes, we assessed the metabolic effects of a mortality-inducing drought on seedlings of *Pinus sylvestris* (Scots Pine), a widespread and important Eurasian species.
- In testing seedlings from provenances with varying water availability, we found divergence over time in the foliar metabolic composition of droughted vs. well-watered individuals, with the former showing increased abundance of aromatic amino acids and decreases in secondary metabolism associated with defence.
- We observed no significant differences amongst provenances in these effects: seedlings from drought-prone areas showed the same foliar metabolic changes under drought as seedlings from moist environments, although morphological effects of drought varied by provenance.
- Overall, our results demonstrate how severe drought may lead to decreases in compounds derived from aromatic amino acids and compromise secondary metabolic pathways related to defences against natural enemies, thereby contributing to the risk of drought-induced mortality in *Pinus sylvestris*.

**Keywords (6-10):** drought, *Pinus sylvestris*, metabolomics, plant defence, seedling, provenance effects, genotype by environment, carbon starvation, hydraulic failure

**Running header (short title):** Drought effects on growth and metabolism of *Pinus sylvestris* seedlings

## INTRODUCTION

Fluctuations in environmental conditions necessitate appropriate plant responses. Despite documentation of widespread forest dieback triggered by drought in all major forested biomes (Allen et al. 2010; Allen et al. 2015), and of elevated mortality under drought in various tree species (Mueller et al. 2005; Breshears et al. 2009; Martinez-Vilalta et al. 2010; Rigling et al. 2013), the underlying mechanisms of drought-induced mortality, species-specific vulnerability and population level resilience are still poorly known. The theoretical framework of physiological mechanisms of tree mortality currently focuses on hydraulic failure, where fluid transport breaks down, and decreasing carbon availability, where metabolic demands may be unmet owing to depletion of non-structural carbohydrates (e.g., McDowell et al. 2008; Galiano et al. 2011; Körner 2015; Adams et al. 2017). Depending on the tree species, these mechanisms need not be mutually exclusive (McDowell et al. 2008; Salmon et al. 2015; Mencuccini et al. 2015). For example, mortality of *Pinus edulis* was found to occur through both hydraulic failure and carbon starvation (Sevanto et al. 2014). Further to these hypotheses, owing to the coupling of xylem and phloem fluid transport, phloem transport limitations under drought have been suggested to contribute to failure to supply the non-structural carbohydrates (NSC) essential to plant metabolism (McDowell & Sevanto 2010; Sevanto et al. 2014). Defining carbon limitations and carbon starvation prior to death has proven hard (e.g., Sala et al. 2012; Sevanto 2014), because little is known about the multiple metabolic pathways via which energy flows to cover the metabolic needs of plants and because of the many ways by which carbon and water limitations can interact in plants.

Owing to high mortality levels during seedling establishment, this stage represents a major bottleneck to recruitment into a population. Seedling establishment has been used as an indicator of the effects of climate change on species assembly in plant communities (Sternberg et al. 1999; Kullman 2002; Lloret et al. 2009). However, despite the heightened mortality rates and increased sensitivity to climate change of this demographic, the mechanism of drought-induced mortality at the stage of seedling establishment is understudied (Lloret et al. 2004, 2009).

Scots pine is widespread across Eurasia, serving as a key timber species and abundant and ecologically important in natural forest stands. Scots pine populations are known to vary ecotypically (Rehfeldt et al. 2002). Latitudinal and longitudinal clines of phenotypic variability in physiological traits have been observed, as well as differences in adaptive plasticity across

provenances (summarised by Semerci et al. 2017). Both phenotypic plasticity and local adaptation could have a role in enabling resilience to drought. Relative to structural and morphological plasticity, metabolism changes over short time-scales and is a more immediate reflection of a plant's response to environmental stressors, including drought. Therefore, a dynamic picture of a plant's response to drought over time can be obtained in metabolic studies, both within and across provenances, which may shed light on how drought-induced mortality is related to particular metabolic pathways. Drought stress is discernible at the level of plant metabolic phenotypes. For example, shifts in carbon metabolism and secondary metabolite synthesis related to water deficit and oxidative stress have been detected in multiple plant species (Zhao et al. 2015, Bowne et al. 2012, Ings et al. 2013, Gargallo-Garriga et al. 2014). Oxidative stress caused by the accumulation of cytotoxic reactive oxygen species (ROS) may be of particular relevance to metabolic activity during mortality-inducing drought, since the extent of oxidative damage is governed by the activity of particular metabolic pathways and the capacity of antioxidant defences to avert an imbalance of ROS (Cruz de Carvalho et al. 2008).

Although the effects of drought on the metabolome of woody plants have been investigated before (e.g., Gargallo-Garriga et al. 2014, 2015; Hamanishi et al. 2015, de Simón et al. 2017), investigation of metabolic changes during drought to the point of mortality has not been carried out. Because the levels of carbon reserves may change and even increase during drought, investigation of the metabolic responses just prior to death are required to determine the sequence of metabolic events leading to mortality (Ryan 2011). Non-targeted metabolomics offers the possibility of capturing a global picture of metabolism, rather than just that related to known metabolic pathways, and it can aid the discovery of novel pathways and interactions amongst pathways (Hall 2006). Both phenotypic plasticity and local adaptation could have a role in enabling resilience to drought and the seedling response is of particular importance, since it represents a bottleneck in terms of higher mortality rates (Castro et al. 2005; Matias et al. 2011; Semerci et al. 2016). Despite increasing knowledge on plasticity and genetic variability of morphological and physiological traits, very little is known about plasticity at the metabolic level. Insights into metabolic changes during seedling drought-induced mortality will help to address this knowledge gap.

Here, we conduct an in-depth comparison of the foliar metabolome of *P. sylvestris* seedlings that were droughted to the point of death versus well-watered, control seedlings. Further, we

compared seedlings from provenances that differ in natural water availability, in order to assess if there is provenance-specific variation in metabolic responses to drought. Additional measurements of biomass and functional traits were taken to determine if changes at the whole-plant level correspond to those observed at the metabolic level. We hypothesise that drought will have major effects on plant metabolism when photosynthesis becomes compromised and carbohydrate availability for plant defence pathways is restricted. We also hypothesise that seedlings from provenances that do not regularly experience intense droughts will show greater metabolic and whole-plant changes under the drought treatment.

## MATERIALS AND METHODS

### *Experimental conditions and sampling*

The experiment was carried out over 5 weeks in July and August of 2015 in a controlled growth chamber at the University of Edinburgh (UK), under constant conditions with diurnal cycles of 16 h light at  $100 \mu\text{mol m}^{-2} \text{s}^{-1}$  and 8 h darkness. The light level was chosen to reduce the rate of water use and consequent dehydration of the plants during the drought, as potted plants would otherwise rapidly undergo hydraulic failure in a manner not reflective of field conditions (Adams et al. 2017). Relative humidity was 65% (day) and 50% (night), with a constant day/night temperature of  $21^{\circ}\text{C}$ . This is the current mean temperature at the southern range limit of this species during July and August (Matías and Jump 2014). The experiment was carried out on 10 month old seedlings, germinated from seeds originating from four provenances spanning a gradient of water availability from wet (Scotland) to intermediate (Austria, Poland) to dry (Spain). Climatic characteristics of the four populations from which seeds were collected are presented in Table 1. Mean annual precipitation was obtained from 1901 to 2015 from the CRU TS3.10 Dataset (Harris et al. 2014). Maximum climatic water deficit values for these sites, which correspond to the summed difference between evapotranspiration and precipitation over the dry season, were obtained from Chave et al. (2014). To give an indication of the degree of water stress expected at the various seed-collection sites, mean soil pF values in the summer month with the maximum mean pF value (July/August) from 1990 to 2014 were obtained from the European Drought Observatory ([edo.jrc.ec.europa.eu/](http://edo.jrc.ec.europa.eu/) on February 2<sup>nd</sup>, 2015). pF expresses the force with which different quantities of water are retained in the soil (Woodruff 1940), with lower values indicating higher water availability for plants. The seeds were collected from open-pollinated trees, with at least 5 maternal parent trees sampled per provenance site. A total

of 1200 seedlings were included in the experiment, with 600 per treatment and 300 per provenance.

Two months prior to instigating drought and control treatments, the seedlings were re-potted into 7 x 7 x 8 cm pots with Levingtons M3 pot and bedding high nutrient mix (Everris, Ipswich, UK). The drought treatment consisted of complete withdrawal of irrigation that resulted in a steep decline in soil water content (Fig. S1). A set of 10 seedlings were droughted one week before the others and were used to indicate when a final sampling point prior to mortality should be carried out. By the end of the first week after the final harvest, 80-90% of seedlings in the drought treatment had died, while mortality had not yet started by the time the final samples had been collected. A one-week, re-watering period was used to confirm that the seedlings were indeed dead. During the experiment, 40 pots across treatments and provenance (5 pots per provenance and treatment) were weighed at 09:00 on days 0, 14, 29 and 36. At the end of the experiment following plant harvesting, the pots were oven dried at 70°C for 48 hours to obtain the dry weight. The gravimetric soil water content ( $\theta_d$ ) (grams of water per gram of oven-dried soil) was calculated as (wet soil weight - dry soil weight) / dry soil weight. Then volumetric water content ( $\theta_{vd}$ ) was calculated as gravimetric soil water content ( $\theta_d$ ) x (bulk density ( $d_b$ ) / density of water ( $d_w$ )).

### *Ecophysiological measurements*

To assess pre-experiment morphological variation amongst provenances, height and crown depth were measured for 15 seedlings per provenance at the start of the experiment. To assess morphological impacts of drought, at each sample point including the start of the experiment ( $t_1=0$ ,  $t_2=11$ ,  $t_3=29$ ), 5 individuals were sampled per provenance per treatment for trait analyses. Fresh leaf weight (FW, g) was measured by separating all needles from the shoot and weighing them. After this, total leaf area (TLA,  $\text{cm}^2$ ) was obtained using scanned images of all leaves and the ImageJ software (Image-J 136b; NIH, Bethesda, Maryland, USA). All needles were saturated in vials of water for 24 hours in order to obtain the turgid weight (TW, g). Dry weights (DW, g) of total needles and stem tissue were obtained after oven-drying for 48 hours at 70 °C. The percent relative water content was then calculated as:  $\text{RWC} = (\text{FW} - \text{DW}) / (\text{TW} - \text{DW}) \times 100$ . Water deficit was calculated as  $\text{WD} = (\text{TW} - \text{FW}) / (\text{TW} - \text{DW})$ . Specific leaf area was calculated as  $\text{SLA} = \text{TLA} / \text{DW}$ .

Entire root systems were washed and arranged so as to avoid overlapping lateral roots on a clear plastic tray with a white background. Images taken with a digital camera were converted to binary and analysed to obtain total root length using RootReader2D plugin (Clark *et al.*, 2013) in ImageJ. Roots were oven dried at 70°C for 48 hours and weighed to measure root biomass (g). Specific root length (SRL, m g<sup>-1</sup>), the ratio of root length to dry mass of roots, was also calculated.

To determine photosynthetic responses to drought, chlorophyll fluorescence from needles was measured by means of a Plant Efficiency Analyzer (Hansatech Instruments Ltd., Norfolk, England). The ratio of variable (F<sub>v</sub>) to maximum fluorescence (F<sub>m</sub>) was taken, since this value has been widely used for assessing plant physiological status and the state of Photosystem II (PSII) (Murchie & Lawson, 2013). F<sub>v</sub>/F<sub>m</sub> was measured for 5 seedlings per provenance at four time points (0, 12, 26 and 33 days). The middle portion of one needle per seedling was placed in the centre of the leaf clip measuring area. Needles were detached and dark adapted in leaf clips supplied with the analyser for 30 min at room temperature. Then the minimum fluorescence (F<sub>o</sub>), maximum fluorescence (F<sub>m</sub>), variable fluorescence (F<sub>v</sub> = F<sub>m</sub> - F<sub>o</sub>), and the ratio F<sub>v</sub>/F<sub>m</sub> were recorded using a saturating intensity pulse for 0.7 s at 80% intensity level of photon flux density (4620 μmol m<sup>-2</sup> s<sup>-1</sup>). The maximal quantum yield at PSII was calculated as  $F_v/F_m = (F_m - F_o)/F_m$ .

#### *Metabolomics sampling, extraction and analysis*

To determine metabolic responses to drought, the foliar tissues of five randomly selected individuals per provenance per treatment were sampled at midday at three time points (0, 11, and 29 days). As the sampling was destructive, these were different individuals than those sampled for trait measurements. Entire shoots were immediately frozen in liquid N<sub>2</sub> to ensure no wounding response was elicited by removing tissue from the plant, and then stored temporarily in frozen aluminium foil on dry ice. Samples were stored in a -80°C freezer until metabolite extraction.

A monophasic extraction method was used to extract metabolites. The solvent comprised acetonitrile, methanol, and HPLC-grade water (2:2:1) and was kept on wet ice to avoid evaporation. Frozen needle tissue was weighed into MK-28 Precellys homogenisation tubes (Stretton Scientific, Stretton, Derbyshire, UK) containing steel beads. The volume of solvent to be added was adjusted according to the fresh tissue weight and 5 % extra acetonitrile and



methanol were added to compensate for estimated differences in relative water content between drought and control. Following the addition of solvent, samples were vortexed and homogenised in a Precellys-24 bead-based homogenizer (Stretton Scientific) at room temperature with two 3 minute pulses of 6800 rpm. The homogenate was transferred to a 1.5 ml microtube (Sarstedt, Nümbrecht, Germany) and half the volume of solvent was used to wash the Precellys tube to ensure that any residual homogenate was transferred. Samples were placed on ice and vortexed for 10 s before the mixture was centrifuged at 19 °C, 14000 rpm (19064 rcf, 10 min). Finally, 400 µl of supernatant were transferred to a new microtube, which was stored at -80°C prior to speed vacuum drying (Wu *et al.*,2008). After centrifugation, equal volumes of the supernatant were transferred into a new plastic tube and dried in a SpeedVac before storage at -80°C. Taking up the samples in 100 µl water / methanol 1:1 with 0.1 % formic acid could not be achieved by vortexing alone and required 20 min of sonication. A quality control (QC) sample was pooled from 10 µl each.

After centrifugation at 15000 rpm (21885 rcf) for 10 min at 4°C (Biofuge), 20 µl per sample were pipetted into a 96-well plate in a controlled randomised order, with QC samples placed equidistantly amongst them. The samples were analysed by UHPLC-MS on a Thermo Scientific Q Exactive mass spectrometer attached to a Thermo Dionex Ultimate 3000 RS system, equipped with a Thermo Hypersil Gold column (100 x 2.1 mm, 1.9 µm particles) (Clark *et al.* 2017). Solvent A was 0.1 % formic acid in water and Solvent B was 0.1 % formic acid in methanol. Liquid chromatography was performed over 14 min at a flow rate of 400 µl/min containing a gradient from A to B from 1 to 8 min. Following analysis of preliminary test samples of aqueous and organic phases, data were collected in positive ion and profile mode,  $m/z$  (mass to charge ratio) 100-1000 Da at 70,000 resolution. Three additional runs of the pooled QC sample were performed, one in positive ion mode at higher resolution (140,000), and two runs using data dependent acquisition; MS/MS fragmentation dependent on the highest 5 signals per MS scan, in positive and negative ion mode. MS data were converted into mzML format using MS Convert, and an R based XCMS/Camera script was used to obtain a first raw intensity matrix, which was imported into MatLab (SimStitch 3.1). Blank filtering was applied using a two-fold sample over blank threshold and peak signal filtering used an 80 % sample filter, applied per group to generate a Sample Filtered Matrix. The dataset was normalised using the probabilistic quotient normalisation (PQN) algorithm to correct for peak intensity differences. Missing values were imputed using a K Nearest Neighbour algorithm (k=5). All

values were then transformed using a generalised logarithm (g-log) to minimise heteroscedasticity in downstream statistical analyses (Parsons et al. 2007; Di Guida et al. 2016).

#### *Statistical and bioinformatics analyses*

In order to assess the effect of drought treatment, provenance and their interaction on ecophysiological traits, we conducted 2-way analyses of variance (ANOVAs). Initial analyses using all time points in the model and time as factor showed that most of the variation in response variables occurred at the third time point and we thus focused on contrasting results for the second and third time points. Where necessary, variables were log or arcsine transformed prior to analysis to improve the normality of model residuals. This applied to: crown percentage of shoot, maximum root length percentage of total root length, leaf RWC and  $F_v/F_m$  data.

Metabolomics data were analysed using a combination of multivariate and univariate statistics. A visual comparison of the QC samples in multivariate space generated by a principal component analysis (PCA) assured us of the technical quality of metabolomics profiles (Fig. S2). The QC samples were then removed from the data for further analyses. We next conducted an additional exploratory PCA to visualise how the overall metabolome varied among provenances, experimental treatments and time points.

To test for the overall response of the metabolome during the experiment, we used regularised multiple analyses of variance (rMANOVA). The combined implementation of multiple analytical approaches has been strongly recommended when analysing metabolomics datasets (Karp et al. 2005; Goodacre et al. 2007; Vinaixa et al. 2012). Multivariate analysis of variance (MANOVA) cannot be used for analysis of high-dimensional data where the number of observations is (much) less than the number of variables. This issue is avoided by using regularized MANOVA (rMANOVA), a multivariate data analysis method that has been specifically developed for analysis of multi-factor untargeted metabolomics data (Engel et al. 2015). rMANOVA can be considered as a MANOVA where a regularized (shrinkage) estimator of the within-group variation is used rather than the sample estimator. Because of this, the method is applicable to data where the number of observations is much smaller than the number of variables. This method is closely related to ANOVA simultaneous component analysis (ASCA), which is a well-known method for analysis of such data. The difference between rMANOVA and ASCA is that rMANOVA tries to better take the correlations between

the observed peak intensities into account, which often makes it a more sensitive method to detect significant differences among groups (Engel *et al.*, 2015). We constructed a full model with factors drought, time point, and provenance, including all possible two-way and three-way interactions. A permutation test with 1000 permutations was used to assess the significance of each factor and interaction in the rMANOVA model (Matlab, R2014b).

ANOVAs, in combination with Benjamini–Hochberg false discovery correction at  $\alpha = 0.05$  (Benjamini & Hochberg 1995), were used for univariate analyses of the data. As provenance did not generally show any significant results on its own or in interaction with other factors, we focused on the effects of sampling time point and drought treatment on metabolite composition of individuals. Specifically, we conducted pairwise comparisons that contrasted different time points and experimental treatments. We first determined the number of metabolites that showed significant differences in abundance between groups, as measured by metabolite peak height.

For metabolite identifications, we focused on metabolites that showed significant differences in abundance between drought and control at time point 3 and had an average absolute  $\log_2$  fold change value greater than 1 in this comparison (i.e. were twice as abundant or twice as rare in droughted versus control seedlings). We focused on this comparison because multivariate and univariate analyses indicated that the greatest differences in metabolic composition were between individuals in drought versus control at time point 3 and because this represents the time point at which individuals in the drought treatment were closest to drought-induced mortality.

Putative metabolite annotations were carried out by matching  $m/z$  values to the Kyoto Encyclopedia of Genes and Genomes (KEGG) (Kanehisa *et al.* 2016), LipidMaps (Fahy *et al.* 2007) and BioCyc (Caspi *et al.* 2016) databases. These annotations were carried out with the software MI-PACK using a 2 ppm (KEGG with molecular formula search up to  $m/z$  620), 3 ppm (BioCyc) or 4 ppm (LipidMaps) error margin, thereby also defining a range of reasonable molecular formulae (Weber & Viant 2010). Since isomers and adducts are included and not always distinguishable, and the number of possible compounds increases over the  $m/z$  range, not all peaks could be assigned with putative metabolite identification. The LC-MS/MS data collected assisted the annotation (level 2 identification; Sumner *et al.* 2007).

To aid in identification of metabolic activity of specific pathways, given incomplete knowledge of individual metabolite identities, the library *mummichog* (Li et al., 2013) from the *MetaboAnalystR* package (Xia et al., 2015) was used. *Mummichog* employs a probabilistic framework to bypass the need for complete prior metabolite identification, by combining knowledge of metabolite spectral features with knowledge of linkages among known metabolites within curated metabolic pathways (based on putative annotations). Given a list of *m/z* ratios, values of log-fold change from control and univariate tests of significance for individual compounds, *mummichog* computes a list of all possible candidate metabolites, including common isotopic derivatives and adducts. If the list of significant features reflects true metabolic activity, the true metabolites should show ‘enrichment’ (i.e., a significant increase in the treatment relative to the control, or vice versa) for particular pathways, while the falsely matched compounds should be distributed more randomly. In practice, *mummichog* calculates a contingency table of enriched metabolites for every known pathway relative to the total number of metabolites in that pathway; it adjusts the Fisher t-test by incorporating the EASE measure of pathway size to increase robustness and draws random permutations of all listed metabolites (including falsely matched compounds) across all pathways to derive a gamma null distribution of P values, which is tested against the observed enrichment list. To identify the metabolic pathways most likely altered immediately prior to death, we employed *mummichog* to test for enrichment in the control against the drought treatment for the last time point of the experiment. We ran *mummichog* in positive analytical mode, assuming a mass accuracy of 3ppm and running 1000 permutations for each run. We considered cut-off probability points varying from 0.05 to 0.0001 for the significance of the gamma distribution and examined which pathways were most consistently identified across all cut-off thresholds (fewer pathways are identified with lower thresholds). Software R version 3.2.2 was used for data analyses (R Core Team, 2015).

## RESULTS

### *Seedling biomass and photochemical capacity*

At the start of the experiment (day 0, time point 1), there were significant differences amongst provenances in height, with the tallest seedlings being from the Austrian provenance, Pernitz (ANOVA, d.f=3, F=3.09, p=0.03). There were marginal differences in SLA, with the highest SLA in seedlings from the Scottish provenance, Rothiemurchus (ANOVA, d.f=3, F=2.81,

p=0.072). Provenances did not differ significantly in any other seedling traits at the start of the experiment ( $p > 0.1$ ). At time point 2 in our study (day 11), most morphological variables were not significantly affected by the drought treatment or provenance. Two exceptions were a significant difference between drought treatment and control for root to shoot ratio (ANOVA, d.f=1,  $F=11.08$ ,  $p=0.002$ ) and significant differences among provenances for maximum quantum yield of PSII,  $F_v/F_m$  (ANOVA, d.f=3,  $F=2.9$ ,  $p=0.04$ ). There were no significant interactions of treatment by provenance at time point 2.

By time point 3 (day 29), greater morphological differences between drought and control treatments were apparent. There was a significant decrease in total, shoot and root dry weight of seedlings under drought, as well as significant provenance interactions with the drought treatment (Table S1; Fig. 1A-C). Seedlings from Rothiemurchus were most affected under drought and showed reduced total, shoot and root dry weight, while the least reduction in biomass was found for the Spanish provenance. Root to shoot ratios significantly decreased under drought, though no provenance interaction effect was found (Fig. 1D). Drought treatment significantly decreased the specific root length and maximum root length by day 29 of the experiment, while no provenance effects were detected (Table S1; Fig. S3). Drought also significantly increased the maximum root length as a fraction of the total root length by day 29 of the experiment (ANOVA, d.f=1,  $F: 11.1$ ,  $p=0.002$ ) (Fig. S4). A significant effect of drought treatment on  $F_v/F_m$  ratios was found after day 26 (Fig. S5). There was also a provenance interaction with drought on  $F_v/F_m$  ratios on day 26.  $F_v/F_m$  decreased most in seedlings from Rothiemurchus and Pernitz and was least reduced in seedlings from Sierra Nevada (Fig. S5).

### *Metabolomics*

Our UHPLC-MS approach generated peak height data for 4640 distinct peaks that putatively represent distinct metabolites with their adducts and isotopes, with peak height representing relative metabolite abundance. As an initial analysis of the data, PCA scores plots were constructed where the samples were coloured according to the levels of the factors of interest (e.g. drought vs control). No separation between the samples was observed except for the interaction between drought and time. The drought time point 3 samples were clearly separated from the other samples (Fig. S6). A principal component analysis-based reduction of the metabolomics data gave two main components that together explained 20% of the variation in the data, while a scree plot showed a sharp reduction in variation explained by subsequent axes.

A visual assessment of these two principal components (Fig. 2A&B) suggests that metabolite composition did not change substantially between time points 1 and 2, but that the leaf metabolome of seedlings in the drought treatment diverged substantially from seedlings in the control treatment at time point 3. For each experimental treatment at each time point, there was substantial overlap amongst provenances in metabolite composition, becoming more pronounced as the experiment progressed (Fig. 2A&B). These visual impressions were confirmed in multivariate analyses of the data by regularised MANOVA (rMANOVA; Engel et al. 2015). This showed a significant effect of drought treatment ( $p < 0.001$ ), time ( $p < 0.001$ ) and their interaction ( $p < 0.001$ ) on leaf metabolic composition of seedlings, while provenance had no significant effects, either on its own ( $p = 0.981$ ) or in interaction with time ( $p = 0.787$ ) or experimental treatment ( $p = 0.987$ ).

Univariate pairwise comparisons of specific time points and treatments showed that a much greater number of metabolites differed significantly in abundance between the drought and control treatments at time point 3 (1138 signals) than at time point 2 (190 signals) (Fig. 2C). Within the drought treatment, seedlings showed a substantial difference in metabolite abundances between time points 2 and 3 (892 signals). Within the control treatment, fewer metabolites changed significantly in abundance between time points 2 and 3 (125 signals) (Fig. S7). The smallest changes in metabolite abundance were observed between seedlings at the start of the experiment and seedlings at time point 2, both in the drought (14 signals) and control (29 signals) treatments (Fig. 2C; Fig S7).

For further scrutiny and identification, we selected signals that showed a significant difference in abundance under drought vs. control at time point 3, and that had a  $\log_2$  fold difference in abundance greater than one (i.e., were twice as abundant or twice as rare under drought vs. control at time point 3). The selection of these metabolites is visualised in a Volcano plot (Fig. 3). Overall, more metabolites experienced a significant and substantial decline under drought ( $\log_2$  fold change  $>1$ ) than a significant and substantial increase (183 vs. 146 signals). Yet, among the metabolites that changed the most under drought vs. control, those that increased showed a greater absolute magnitude of change than those that decreased (Fig. 3). For example, among the top 25 signals in terms of absolute magnitude of change, 23 of those increased under drought vs. control, while only two decreased.

We were able to obtain putative annotations for a minority of the signals that showed significant decreases or increases in abundance under drought vs. control. Of the 183 peaks that were significantly less abundant under drought and show a  $\log_2$  fold change less than -1 (i.e. are at least twice as rare on average in drought vs. control at time point 3), only 29 were matched in the KEGG, LIPDMAPS or BIOCYC databases. Of the 49 signals with the greatest decrease under drought relative to control, only 10 have putative annotations in these databases (Table 2; see Table S2 for full list). Of the 146 peaks that are significantly more abundant in drought vs. control and show a  $\log_2$  fold change of at least 1 (i.e. are at least twice as common on average in drought vs. control), 98 have no putative annotations in the KEGG, LIPIDMAPS or BIOCYC databases. Of the 22 compounds with the greatest increase under drought relative to control in terms of fold-changes, only 10 have putative annotations (Table 3; see Table S3 for full list). The metabolites that were more abundant under drought were mainly aromatic amino acids, which may signify up-regulation of aromatic amino acid biosynthesis via the shikimate pathway or protein degradation under drought stress. A number of upregulated signals coeluting at 4.08 min are likely breakdown products of only one metabolite, tryptophan (manual interpretation).

Pathway analysis with *mummichog* identified the following pathways as the primary sites of metabolic enrichment immediately prior to death: aminoacyl-tRNA, phenylalanine (Phe), vitamin B6, phenylpropanoid, isoquinoline alkaloid, arginine and proline (Pro), Phe and Tyrosine (Tyr) and Tryptophan (Trp) biosynthesis (Table S4). Interestingly, the amino acids synthesised via the shikimate pathway (Tyr, Trp and Phe) clearly increased during drought, while several of those in the downstream pathways from shikimate decreased, although the downstream response was compound-specific. Many of the significant decreases during drought were seen for mevalonate and compounds in the mevalonate/MEP-DOXP (or non-mevalonate) pathways, such as carotenoids, sterols, ecdysones, terpenoids and glycosides with a second active group of the same type.

## DISCUSSION

This study investigated the foliar metabolic responses of *Pinus sylvestris* seedlings to severe drought, which eventually led to drought-induced mortality, and to ascertain whether ecotypic adaptation in the metabolic responses to drought would be discernible amongst provenances. Pronounced metabolic effects of drought were found, potentially related to a decoupling of

carbon assimilation and secondary metabolism as photosynthesis became compromised. This manifested through an increase in free amino acids and a decrease in secondary metabolite production via the shikimate and mevalonate/MEP-DOXP pathways. Our second hypothesis was not supported, as we found no significant foliar metabolic differences amongst provenances in the response to drought. The latter is particularly interesting in light of the fact that we found morphological responses to drought to vary by provenance. Seedlings from the driest provenance were least affected morphologically by drought.

#### *Seedling foliar metabolome response to drought*

There were significant differences in the metabolite composition of the control and drought treatment groups, with a clear divergence in metabolic profiles following four weeks of drought stress after which mortality rapidly ensued (Fig. 2). Free amino acids (Tyr, Trp and Pro; Table 3) produced during primary metabolism were identified by ANOVA and rMANOVA. Tyr and Trp increased with the highest fold change under the drought treatment. Pathway analysis with *mummichog* highlighted the aminoacyl t-RNA pathway as enriched, owing to the abundance of free amino acids. However, the inference of increased t-RNA synthesis based solely on the presence of free amino acids seems untenable, especially given that drought stress conditions would presumably have impeded protein synthesis. Besides the already mentioned aromatic amino acids Tyr and Trp, significant increases were also found in several other amino acid metabolic pathways (i.e., phenylalanine, proline, glutamine, valine, leucine, isoleucine, arginine, histidine), leaving only the metabolic pathways of cysteine, methionine, glycine, serine and threonine unaffected. The shikimate pathway involved in aromatic amino acid biosynthesis was identified as a dominant pathway by *mummichog*. Plants synthesise aromatic amino acids, such as Tyr and Trp, via the metabolically costly 7-step shikimate pathway, to which over 30% of photosynthetically derived carbon can be directed (Maeda & Dudareva 2012). Secondary compounds involved in plant defence and abiotic stress responses can be derived from shikimate. It is conceivable that, rather than biosynthesis via the shikimate pathway, protein degradation accounts for the increasing levels of free amino acids detected. The higher abundance of aromatic amino acids relative to other amino acids however seems to indicate that the shikimate pathway is involved. Nevertheless, a concomitant increase in shikimate and chorismate intermediates was not found, which has been used to confirm involvement of the shikimate pathway in accumulation of free aromatic amino acids (Tyr, Trp and Phe) in drought sensitive wheat leaves (Michaletti *et al.* 2017). Pathway analysis



furthermore identified the mevalonate pathways in secondary metabolism as most affected by drought, with decreases in key metabolites involved with plant defence.

Tyrosine (Tyr) was the metabolite that showed the strongest increase under the drought treatment. Tyr hyperaccumulation in young shade leaves of *Inga umbellifera* has been linked to decreased insect larval performance, thus representing a rare example of an amino acid functioning as a defensive compound (Lokvam *et al.*, 2006). Among the annotated metabolites, tryptophan (Trp) showed the third greatest increase under drought versus control. Trp biosynthetic enzymes have been shown to be up-regulated in response to oxidative stress treatment in *Arabidopsis* (Zhao *et al.*, 1998). The increase in aromatic and non-aromatic amino acids, some of which are key precursors of important defence and antioxidant pathways, occurred while several defence compounds decreased (Table S4). Significant decreases also occurred for phosphomevalonate and other downstream compounds that have similar biological roles.

For coniferous species, carbon-based secondary metabolites, terpenoids and phenolic compounds are expressed constitutively and are inducible to high concentrations and provide effective defence against many pests and pathogens (Keeling & Bohlmann 2006). Decreases of secondary metabolites were detected at 29 days of drought, by ANOVA and rMANOVA. This decrease in metabolites involved in defence during the last phase of drought may reflect a decreased capacity to employ the mevalonate pathways at a stage during drought where photosynthesis and growth were already strongly impacted, and carbohydrate availability was limited.

Increased free amino acids under stress may also be the result of protein breakdown, rather than *de novo* biosynthesis of amino acids. For example, branched chain amino acids and aromatic amino acids (Tyr and Phe) have been shown to increase under osmotic stress via protein degradation in *Arabidopsis thaliana*, with a higher fold change than other amino acids; this increase was potentially owing to lower basal levels of branched chain amino acids and aromatic amino acids prior to stress induction, or slower catabolism subsequently (Huang & Jander 2017). In *Arabidopsis*, accumulation of aromatic amino acids occurs more rapidly than branched chain amino acids (Fàbregas and Fernie 2019). It is possible that the accumulation of free aromatic amino acids has a protective role under conditions of elevated oxidative stress, by scavenging free radicals (Stadtman & Levine 2003). Indeed, 2-phenylacetamide was ranked

as the second most elevated metabolite under drought and is a product of oxidation of the aromatic amino acid phenylalanine. However, the pool size changes in amino acids may simply represent an artefact of protein breakdown coupled with differential rates of amino acid catabolism. Under conditions of carbon starvation, catabolism of amino acids would provide a source of energy. The degradation pathways (catabolism) of aromatic amino acids Phe and Trp are not well-documented in plants (Hildebrandt et al. 2015). A comprehensive study of amino acid contents that included low abundance amino acids (such as aromatic ones) showed that the amount of aromatic amino acids produced by protein degradation was sufficient for secondary metabolite production, with the exception of Trp under certain stress conditions (Hildebrandt et al. 2018). Proteolysis can account for the accumulation of amino acids under osmotic stress and fast catabolism of Lys as well as branched chain amino acids has been shown to be induced to provide alternative respiratory substrate during drought stress (Araujo et al. 2011; Batista-Silva et al. 2019; Pires et al. 2016). Thus turnover of amino acids may be explained in terms of proteolysis and tightly regulated amino acid metabolism.

One of the compounds with the largest increase under drought was proline. Proline is an amino acid that functions as a compatible solute, carrying no net charge at physiological pH, by raising osmotic pressure in the cytoplasm and stabilising proteins and cellular membranes (Szabados & Savoure 2010). Additionally, proline has been indicated to exhibit antioxidant capacity in free radical scavenging (Smirnoff & Cumbes 1989). Whereas aromatic amino acid accumulation under osmotic stress was linked to protein breakdown in *Arabidopsis*, increased biosynthesis accounted for elevated levels of proline (Huang & Jander 2017). Indeed, proline is biosynthesised in a number of plant species under drought and has been shown to exhibit osmoprotective and antioxidant capacities (Hayat *et al.*, 2012). Furthermore, it is suggested that proline accumulation under stress may have a role in signalling as well as maintenance of NAD(P)/NAD(P)H ratios that enable metabolic pathways to function in generating secondary metabolites (Hare & Cress 1997). Biosynthesis of proline under water deficit occurs in an abscisic acid-dependent manner and can also be influenced by sugar availability, indicating a regulatory mechanism to restrict proline accumulation when carbohydrate status is low (Rook *et al.*, 2001, Verslues & Bray 2006).

A number of important, significant peaks determined by univariate and multivariate statistics were not identifiable through interrogation of metabolic databases, possibly owing to taxon specificity and the poor development of metabolite databases for non-model plant species. This

is in addition to adducts and breakdown products often not covered by common databases. This finding highlights the difficulty of current metabolomics approaches, at least for conifers, and suggests that insufficient understanding of the metabolic pathways affected by drought is currently hindering our understanding of the nature of carbon limitations prior to drought-induced mortality.

#### *Provenance variation in the seedling drought response*

Total seedling biomass was reduced under drought treatment and this reduction varied significantly across provenances. Seedlings from the wettest provenance, Rothiemurchus (Scotland), were most affected under drought in terms of biomass reduction, while seedlings from the driest provenance, Sierra Nevada (Spain), were least affected. Under drought, total leaf area was significantly reduced, but there was no observed needle abscission. Rather, the drought is thought to have inhibited needle growth. Needles also showed a reduction in the maximal efficiency of photosystem II ( $F_v/F_m$ ) by day 26 of the drought treatment, indicating that photosynthesis was compromised, with the wettest provenance being most affected.  $F_v/F_m$  declines when water stress becomes severe (Epron & Dreyer 1992; Iijima *et al.*, 2006; Ditmarova *et al.*, 2010; Way *et al.*, 2013). Potentially this could be a result of decoupling between photosynthesis and secondary metabolism, with C limitation once assimilation is not occurring owing to stomatal closure in this isohydric species.

A limitation of this study is the lack of information on how mortality rate varies by provenance. Matias and Jump (2014) found that mortality rate of *Pinus sylvestris* under drought is strongly affected by temperature, with the southern Spanish provenance showing higher survival probabilities than the northern Finnish provenance under drought at the current temperature regime of the southern range limit. This effect may not be significant in our experiment owing to the shorter timeframe (5 weeks compared with 19 weeks), since Matias and Jump found no significant effect of temperature on provenances until after 5 weeks of treatment. Survival is vital in determining seedling recruitment at the population level, thus if mortality rate is relatively constant among provenances over short intense drought episodes, then the superior drought response of the Spanish provenance in terms of morphological and physiological traits would not have adaptive significance.

In contrast, metabolic profiles were not found to differ significantly by provenance, a result robust to changes in the signal filter threshold employed. Previous studies on intraspecific variation in conifers have found evidence for a stronger environmental than genetic signal on metabolomes of developing xylem in *Pseudotsuga menziesii* (Robinson et al. 2007). However, differences in foliar metabolomes of *Pinus pinaster* were found to be strongly related to the aridity of the provenance site of origin, suggesting local adaptation (Meijon et al. 2016). Also, Du et al. (2015) found that *Pseudotsuga menziesii* seedlings showed a provenance specific drought response, with the drier provenance increasing aromatic amino acids. In this study, the metabolic phenotypes of *P. sylvestris* seedlings do not appear to be population specific or to show local adaptation, or at least such differences did not have a sufficiently large effect size to be statistically detectable. *Pinus sylvestris* is a widespread species with high plasticity evident across European populations in traits related to physiology, phenology and morphology (Olekysn et al. 1998, 2000; Semerci et al. 2017), but the findings of this study suggest a limitation of foliar metabolic plasticity to react to extreme mortality-inducing drought episodes. The relatively long mean leaf life span of *Pinus* species (over 3 years) may account for the discrepancy between the drought response in foliar metabolic profiles and other measured leaf traits. In evolutionary terms, leaf longevity has a pivotal role in the leaf economic spectrum, with trade-offs between persistence and productivity constraining both morphological and biochemical leaf traits (Warren and Adams 2000; Onoda et al. 2017). Plasticity in needle longevity is known for *Pinus sylvestris* (Pensa and Jalkanen 2005). Juvenile needles in three Mediterranean pine species were found to exhibit a leaf strategy to maximise carbon gain and the transition to adult needles reflected changes in traits according to the drought stress tolerance of each species (Kuusk et al. 2017). Foliar integration of traits has been found to be a function of ontogeny; canalisation of leaf traits are found in plants of reproductive age, whereas in juvenile plants weaker correlations among functional leaf traits indicate lower foliar integration and higher phenotypic plasticity (Damian et al. 2016). However, the selection of leaf traits in this study did not include biochemical or metabolic functional traits. Metabolically divergent drought responses in mature trees may be more likely than for seedlings with the same leaf age, which all show a propensity to maximise carbon gain to the detriment of environmental stress tolerance; thus, canalisation of leaf traits may have affected metabolic profiles.

## Conclusions

There was a strong impact of drought at the metabolic level in *P. sylvestris* needles, with the effect becoming very apparent just before seedling mortality. In response to drought, we found increases in many free amino acids and decreased concentrations of secondary metabolites in the mevalonate/MEP-DOXP pathways. The identification of decreases in compounds derived from aromatic amino acids and secondary metabolic pathways related to plant defence shows that these metabolically costly pathways are down-regulated under drought stress, revealing the incapacity of severely droughted seedlings to produce defensive compounds against biotic stress. However, proline, a compound important for osmoregulation, plant signalling and antioxidant defence, was strongly upregulated, suggesting that its high concentration was important in the latest phases of survival prior to death. While seedling biomass and photochemical efficiency were found to be most strongly reduced by drought for Rothiemurchus, the wettest provenance, there was a lack of provenance effects on metabolite abundances. Overall, our findings indicate that important metabolite changes under drought were centred around the shikimate and the mevalonate/MEP-DOXP pathways. It also demonstrates that a large number of unknown compounds were affected by drought and are therefore of interest for future research.

## Acknowledgements

This research was partly funded via Natural Environment Research Council (NERC) grant NE/I011749/1 to MM and via a NERC PhD studentship to SM. A grant from the NERC Biomolecular Analysis Facility (NBAF868) made the metabolomics study possible.

## Author Contribution

S.M and K.G.D. designed and executed the experiment; S.M measured plant traits and extracted the metabolites; U.S and J.E led the metabolomics analyses; K.G.D and M.M conducted additional analyses; S.M wrote the manuscript; and all authors contributed to revisions.

## REFERENCES

Allen CD, Macalady AK, Chenchouni H, Bachelet D, McDowell N, Vennetier M, Kitzberger T, Rigling A, Breshears DD, Hogg ET, Gonzalez P (2010) A global overview of

- drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest Ecology and Management* 259: 660-684.
- Adams H et al. (2017) A multi-species synthesis of physiological mechanisms in drought-induced tree mortality. *Nature Ecology and Evolution*, 1: 1285-1291.
- Allen CD, Breshears DD, McDowell NG (2015) On underestimation of global vulnerability to tree mortality and forest die-off from hotter drought in the Anthropocene. *Ecosphere* 6 1-55.
- Anderegg WR, Hicke JA, Fisher RA, Allen CD, Aukema J, Bentz B, Hood S, Lichstein JW, Macalady AK, McDowell NG and Pan Y (2015) Tree mortality from drought, insects, and their interactions in a changing climate. *New Phytologist* 208: 674-683.
- Araújo, WL, Tohge, T, Ishizaki, K, Leaver, CJ, Fernie, AR(2011). Protein degradation—An alternative respiratory substrate for stressed plants. *Trends in Plant Science*, 16, 489–498. <https://doi.org/10.1016/j.tplants.2011.05.008>
- Batista-Silva, W., Heinemann, B., Rugen, N., Nunes-Nesi, A., Araújo, W. L., Braun, H. P., & Hildebrandt, T. M. (2019). The role of amino acid metabolism during abiotic stress release. *Plant, cell & environment*.
- Benjamini Y, Hochberg Y. (1995) Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc B* 57: 289–300.
- Bowne JB, Erwin TA, Juttner J, Schnurbusch T, Langridge P, Bacic A, Roessner U (2012) Drought responses of leaf tissues from wheat cultivars of differing drought tolerance at the metabolite level. *Molecular Plant* 5: 418–429.
- Breshears DD, Cobb NS, Rich PM, Price KP, Allen CD, Balice RG *et al.* (2005) Regional vegetation die-off in response to global-change-type drought. *Proceedings of the National Academy of Sciences, USA* 102: 15144–8.
- Caspi R., Billington R., Ferrer L., Foerster H., Fulcher C.A., Keseler I.M., Kothari A., Krummenacker M., Latendresse M., Mueller L.A., Ong Q., Paley S., Subhraveti P., Weaver D.S., Karp P.D(2016) The MetaCyc Database of metabolic pathways and enzymes and the BioCyc collection of Pathway/Genome Databases. *Nucleic Acids Research* 44(D1):D471-80.
- Chave J, Réjou-Méchain M, Búrquez A, Chidumayo E, Colgan MS, Delitti WBC *et al.* (2014) Improved pantropical allometric models to estimate the above ground biomass of tropical forests. *Global Change Biology*, 20:3177-3190.
- Clark, M.S., Sommer, U., Sihra, J.K., Thorne, M.A., Morley, S.A., King, M., Viant, M.R. and Peck, L.S., (2017) Biodiversity in marine invertebrate responses to acute warming

693 revealed by a comparative multi-omics approach. *Global Change Biology*, 23(1),  
 694 pp.318-330.

695 Clark RT, Famoso AN, Zhao K, Shaff JE, Craft EJ, Bustamante CD *et al.* (2013) High-  
 696 throughput two-dimensional root system phenotyping platform facilitates genetic  
 697 analysis of root growth and development. *Plant Cell & Environment* 36: 454–466.

698 Cruz de Carvalho MH. (2008) Drought stress and reactive oxygen species. *Plant Signaling &*  
 699 *Behavior* 3: 156–165.

700 Damián X, Fornoni J, Domínguez CA and Boege K, (2018) Ontogenetic changes in the  
 701 phenotypic integration and modularity of leaf functional traits. *Functional ecology*,  
 702 32:234-246.

703 de Simón, B. F., Sanz, M., Cervera, M. T., Pinto, E., Aranda, I., & Cadahía, E. (2017). Leaf  
 704 metabolic response to water deficit in *Pinus pinaster* Ait. relies upon ontogeny and  
 705 genotype. *Environmental and experimental botany*, 140, 41-55.

706 Di Guida R, Engel J, Allwood JW, Weber RJ, Jones MR, Sommer U, Viant MR, Dunn WB  
 707 (2016) Non-targeted UHPLC-MS metabolomic data processing methods: A  
 708 comparative investigation of normalisation, missing value imputation, transformation  
 709 and scaling. *Metabolomics* 12: 1-14.

710 Ditmarova L, Kurjak D, Palmroth S, Kmet J, Strelcova K. (2010) Physiological responses of  
 711 Norway spruce (*Picea abies* (L.) Karst) seedlings to drought stress. *Tree Physiology*  
 712 30: 205–213.

713 Dixon RA (2001) Natural products and plant disease resistance. *Nature* 411: 843-847.

714 Du B, Jansen K, Kleiber A, Eiblmeier M, Kammerer B, Ensminger I, Gessler A, Rennenberg  
 715 H, Kreuzwieser J. (2015) A coastal and an interior Douglas fir provenance exhibit  
 716 different metabolic strategies to deal with drought stress. *Tree Physiology* 36: 148–163.

717 Engel J, Blanchet L, Bloemen B, Van den Heuvel LP, Engelke, UHF, Wevers RA, Buydens  
 718 LMC(2015) Regularized MANOVA (rMANOVA) in untargeted metabolomics.  
 719 *Analytica Chimica Acta* 899: 1-12.

720 Epron D, Dreyer E (1992) Effects of severe dehydration on leaf photosynthesis in *Quercus*  
 721 *petraea* (Matt) Liebl – photosystem-II efficiency, photochemical and non-  
 722 photochemical fluorescence quenching and electrolyte leakage. *Tree Physiology* 10:  
 723 273–284.

724 Fàbregas N and Fernie AR, (2019). The metabolic response to drought. *Journal of*  
 725 *Experimental Botany*, 70:1077-1085.

726 Facchini PJ (2001) Alkaloid biosynthesis in plants: biochemistry, cell biology, molecular  
 727 regulation, and metabolic engineering applications. *Annual Review of Plant Biology*  
 728 52: 29-66.

729 Fahy E, Sud M, Cotter D, Subramaniam S (2007) LIPID MAPS online tools for lipid research.  
 730 *Nucleic Acids Res* 35:W606–W612.

731 Galiano L, Martínez-Vilalta J, Lloret F (2011) Carbon reserves and canopy defoliation  
 732 determine the recovery of Scots pine 4 yr after a drought episode. *New Phytologist* 190:  
 733 750-759.

734 Gargallo-Garriga A, Sardans J, Pérez-Trujillo M, Rivas-Ubach A, Oravec M, Vecerova K,  
 735 Urban O, Jentsch A, Kreyling J, Beierkuhnlein C, Parella T, Peñuelas J (2014)  
 736 Opposite metabolic responses of shoots and roots to drought. *Scientific Reports*, 4:  
 737 6829, 10.1038/srep06829

738 Goodacre R, Broadhurst D, Smilde AK, Kristal BS, Baker DJ, et al. (2007) Proposed minimum  
 739 reporting standards for data analysis in metabolomics. *Metabolomics* 3: 231–241.

740 Hall RD (2006) Plant metabolomics: from holistic hope, to hype, to hot topic. *New Phytologist*  
 741 169: 453–468.

742 Hamanishi, E. T., Barchet, G. L., Dauwe, R., Mansfield, S. D., & Campbell, M. M. (2015).  
 743 Poplar trees reconfigure the transcriptome and metabolome in response to drought in a  
 744 genotype-and time-of-day-dependent manner. *BMC genomics*, 16(1), 329.

745 Hare PD, Cress WA (1997) Metabolic implications of stress-induced proline accumulation in  
 746 plants. *Plant Growth Regulation* 21: 79-102.

747 Harris I, Jones PD, Osborn TJ, Lister DH (2014) Updated high-resolution grids of monthly  
 748 climatic observations - the CRU TS3.10 Dataset. *International Journal of*  
 749 *Climatology* 34, 623-642.

750 Hayat S, Hayat Q, Alyemeni MN, Wani AS, Pichtel J, Ahmad A(2012) Role of proline under  
 751 changing environments: A review. *Plant Sign & Behaviour* 7: 1456–1466.

752 Hildebrandt TM, Nunes Nesi A, Araujo WL, Braun HP (2015) Amino acid catabolism in  
 753 plants. *Molecular Plant* 8: 1563 –1579.

754 Hildebrandt, T. M. (2018). Synthesis versus degradation: directions of amino acid metabolism  
 755 during *Arabidopsis* abiotic stress response. *Plant molecular biology*, 98(1-2), 121-135.

756 Huang T, Jander G(2017) Absciscic acid-regulated protein degradation causes osmotic stress-  
 757 induced accumulation of branched-chain amino acids in *Arabidopsis thaliana*. *Planta*  
 758 246: 737–747.



759 Iijima H, Shibuya M, Saito H, Takahashi K (2006) The water relation of seedlings of *Picea*  
 760 *jezoensis* on fallen logs. Canadian Journal of Forest Research 36: 664–670.

761 Ings J, Mu LA, Robson PR, Bosch M (2013) Physiological and growth responses to water  
 762 deficit in the bioenergy crop *Miscanthus x giganteus*. Frontiers in Plant Science 4: 468.

763 Kanehisa M, Sato Y, Kawashima M, Furumichi M, Tanabe M (2016) KEGG as a reference  
 764 resource for gene and protein annotation. Nucleic Acids Research 44: D457-D462.

765 Karp NA et al (2005) Impact of replicate types on proteomic expression analysis. Journal of  
 766 Proteome Research 4:1867–1871.

767 Keeling CI, Bohlmann J (2006) Genes, enzymes and chemicals of terpenoid diversity in the  
 768 constitutive and induced defence of conifers against insects and pathogens. New  
 769 Phytologist 170: 657–675.

770 Körner C (2015) Paradigm shift in plant growth control. Current Opinion in Plant  
 771 Biology 25: 107–114.

772 Kullman L (2002) Rapid recent range-margin rise of tree and shrub species in the Swedish  
 773 Scandes. Journal of Ecology. 90: 68-77Li S, Park Y, Duraisingham S, Strobel FH, Khan  
 774 N, Soltow QA, Jones DP, Pulendran B( 2013) Predicting network activity from high  
 775 throughput metabolomics. PLoS Computational Biology 9: e1003123.

776 Kuusk V, Niinemets Ü. and Valladares F, (2017) A major trade-off between structural and  
 777 photosynthetic investments operative across plant and needle ages in three  
 778 Mediterranean pines. Tree physiology, 38:543-557.

779 Lloret F, Peñuelas J, Estiarte M (2004) Experimental evidence of reduced diversity of seedlings  
 780 due to climate modification in a Mediterranean-type community. Global Change  
 781 Biology. 10: 248-258.

782 Lloret F, Peñuelas J, Prieto P, Llorens L, Estiarte M (2009) Plant community changes induced  
 783 by experimental climate change: Seedling and adult species composition. Perspectives  
 784 in Plant Ecology, Evolution and Systematics. 11: 53-63.

785 Lokvam J, Brenes-Arguedas T, Lee JS, Coley PD, Kursar TA. (2006) Allelochemic function  
 786 for a primary metabolite: the case of L-tyrosine hyper-production in *Inga umbellifera*  
 787 (Fabaceae). American Journal of Botany, 93: 1109-1115.

788 Maeda H, Dudareva N (2012) The Shikimate pathway and aromatic amino acid biosynthesis  
 789 in plants. Annual Review of Plant Biology 63: 73–105.

790 Martinez-Vilalta J, Mencuccini M, Vayreda J, Retana J (2010) Interspecific variation in  
 791 functional traits, not climatic differences among species ranges, determines

792 demographic rates across 44 temperate and Mediterranean tree species. *Journal of*  
793 *Ecology* 98: 1462–1475.

794 Matías L, Jump AS (2014) Impacts of predicted climate change on recruitment at the  
795 geographical limits of Scots pine. *Journal of Experimental Botany* 65: 299–310.

796 McDowell N, Pockman WT, Allen CD, Breshears DD, Cobb N, Kolb T *et al.* (2008)  
797 Mechanisms of plant survival and mortality during drought: why do some plants  
798 survive while others succumb to drought? *New Phytologist* 178: 719–39.

799 McDowell NG, Sevanto S (2010) The mechanisms of carbon starvation: how, when, or does it  
800 even occur at all? *New Phytologist* 186: 264–266.

801 McDowell NG, Beerling DJ, Breshears DD, Fisher RA, Raffa KF, Stitt M (2011) The  
802 interdependence of mechanisms underlying climate-driven vegetation mortality.  
803 *Trends in Ecology and Evolution* 26:523–532.

804 Meijon M, Feito I, Oravec M, Delatorre C, Weckwerth W, Majada J, Valledor L (2016)  
805 Exploring natural variation of *Pinus pinaster* Aiton using metabolomics: Is it possible  
806 to identify the region of origin of a pine from its metabolites? *Molecular Ecology* 25:  
807 959–976.

808 Mencuccini M, Minunno F, Salmon Y, Martínez-Vilalta J, Hölttä T, (2015) Coordination of  
809 physiological traits involved in drought-induced mortality. *New Phytologist*, 208: 396-  
810 409.

811 Michaletti A, Reza Naghavi M, Toorchi M, Zolla L & Rinalducci S (2018) Metabolomics and  
812 proteomics reveal drought-stress responses of leaf tissues from spring-wheat. *Nature*  
813 *Scientific Reports* 8: 5710.

814 Mueller RC, Scudder CM, Porter ME, Trotter RT, Gehring CA, Whitham TG (2005)  
815 Differential tree mortality in response to severe drought: evidence for long-term  
816 vegetation shifts. *Journal of Ecology* 93: 1085–1093.

817 Murchie EH, Lawson T (2013) Chlorophyll fluorescence analysis: a guide to good practice and  
818 understanding some new applications. *Journal of Experimental Botany* 64: 3983–3998.

819 Oleksyn J, Zytowskiak R, Karolewski P, Reich PB, Tjoelker MG (2000) Genetic and  
820 environmental control of seasonal carbohydrate dynamics in trees of diverse *Pinus*  
821 *sylvestris* populations. *Tree Physiology* 20: 837–847.

822 Oleksyn J, Tjoelker MG, Reich PB (1998) Adaptation to changing environment in Scots pine  
823 populations across a latitudinal gradient. *Silva Fennica* 32: 129–140.

824 Onoda Y, Wright IJ, Evans JR, Hikosaka K, Kitajima K, Niinemets Ü, Poorter H, Tosens T  
825 and Westoby M (2017) Physiological and structural tradeoffs underlying the leaf  
826 economics spectrum. *New Phytologist*, 214:1447-1463.

827 Parsons HM, Ludwig C, Günther UL, Viant MR (2007) Improved classification accuracy in 1-  
828 and 2-dimensional NMR metabolomics data using the variance stabilising generalised  
829 logarithm transformation. *BMC Bioinformatics* 8: 234.

830 Pensa M and Jalkanen R (2005) Variation in needle longevity is related to needle-fascicle  
831 production rate in *Pinus sylvestris*. *Tree physiology*, 25:1265-1271.

832 Pires MV, Pereira Júnior AA, Medeiros DB, Daloso DM, Pham PA, Barros KA, Engqvist MK,  
833 Florian A, Krahner I, Maurino VG, et al. The influence of alternative pathways of  
834 respiration that utilize branched-chain amino acids following water shortage in  
835 *Arabidopsis*. *Plant Cell Environ.* 2016, 39, 1304–1319.

836 R Core Team. 2015. *R: A language and environment for statistical computing*. R Foundation  
837 for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.

838 Rehfeldt GE, Tchebakova NM, Perfenova YI, Wykoff WR, Kuzmina NA & Milyutin LI (2002)  
839 Intraspecific responses to climate in *Pinus sylvestris*. *Global Change Biology* 8: 919-  
840 929.

841 Rigling A, Bigler C, Eilmann B, Feldmeyer-Christe E, Gimmi U, Ginzler C *et al.* (2013)  
842 Driving factors of a vegetation shift from Scots pine to pubescent oak in dry Alpine  
843 forests. *Global Change Biology* 19: 229–240.

844 Robinson AR, Ukrainetz NK, Kang KY, Mansfield SD (2007) Metabolite profiling of Douglas-  
845 fir (*Pseudotsuga menziesii*) field trials reveals strong environmental and weak genetic  
846 variation. *New Phytologist* 174: 762–73.

847 Rook F, Corke F, Card R, Munz G, Smith C, Bevan MW (2001) Impaired sucrose-induction  
848 mutants reveal the modulation of sugar-induced starch biosynthetic gene expression by  
849 abscisic acid signalling. *The Plant Journal* 26: 421-433.

850 Ryan MG (2011) Tree responses to drought. *Tree Physiology* 31, 237–239.

851 Sala A, Woodruff DR, Meinzer FC(2012) Carbon dynamics in trees: feast or famine? *Tree*  
852 *Physiology* 32:764–75.

853 Salmon Y, Torres-Ruiz JL, Poyatos R, Martinez-Vilalta J, Meir P, Cochard H, Mencuccini M  
854 (2015) Balancing the risks of hydraulic failure and carbon starvation: a twig scale  
855 analysis in declining Scots pine. *Plant, Cell & Environment* 38: 2575-2588.

856 Semerci A, Semerci H, Çalışkan B, Çiçek N, Ekmekçi Y, Mencuccini M (2017) Morphological  
857 and physiological responses to drought stress of European provenances of Scots pine.  
858 European Journal of Forest Research 136: 1-14.

859 Sevanto S, McDowell NG, Dickman LT, Pangle R, Pockman WT (2014) How do trees die? A  
860 test of the hydraulic failure and carbon starvation hypotheses. Plant, Cell &  
861 Environment 37: 153-161.

862 Smirnoff N, Cumbes J (1989) Hydroxyl radical scavenging activity of compatible solutes.  
863 Phytochemistry 28: 1057–1060.

864 Stadtman ER, Levine RL (2003) Free radical-mediated oxidation of free amino acids and  
865 amino acid residues in proteins. Amino Acids 25: 207-218.

866 Sternberg, M., Brown, V.K., Masters, G.J. et al. Plant Ecology (1999) 143: 29.

867 Sumner LW, Amberg A, Barrett D, Beale MH, Beger R, Daykin CA *et al.* (2007) Proposed  
868 minimum reporting standards for chemical analysis. Metabolomics 3:211–221.

869 Szabados L, Savoure A (2010) Proline: a multifunctional amino acid. Trends Plant Science 15:  
870 89-97.

871 Verslues PE, Bray E (2006) Role of abscisic acid (ABA) and Arabidopsis thaliana ABA-  
872 insensitive loci in low water potential-induced ABA and proline accumulation. Journal  
873 of Experimental Botany 57: 201–12.

874 Vinaixa M, Samino S, Saez I, Duran J, Guinovart JJ, Yanes O (2012) A guideline to univariate  
875 statistical analysis for LC/MS-based untargeted metabolomics-derived  
876 data. Metabolites. 2:775–95.

877 Warren CR and Adams MA (2000) Trade-offs between the persistence of foliage and  
878 productivity in two Pinus species. Oecologia, 124:487-494.

879 Way DA, Crawley C, Sage RF (2013) A hot and dry future: Warming effects on boreal tree  
880 drought tolerance. Tree Physiology 33: 1003–1005.

881 Weber, R. J. M., & Viant, M. R. (2010). MI-Pack: Increased confidence of metabolite  
882 identification in mass spectra by integrating accurate masses and metabolic pathways.  
883 Chemometrics and Intelligent Laboratory Systems, 104: 75–82.

884 Woodruff CM (1940) Soil moisture and plant growth in relation to pF. Soil Science Society  
885 America Proceedings 5: 36-41.

886 Wu Q-S, Xia R-X, Zou Y-N (2008) Improved soil structure and citrus growth after inoculation  
887 with three arbuscular mycorrhizal fungi under drought stress. European Journal of Soil  
888 Biology 44: 122–128.

889 Xia J, Sinelnikov I, Han B, Wishart DS (2015) MetaboAnalyst 3.0 - making metabolomics  
890 more meaningful. *Nucleic Acids Research* 43: W251-257.

891 Zhao J, Last RL, Williams CC( 1998) Induction of Arabidopsis tryptophan pathway enzymes  
892 and camalexin by amino acid starvation, oxidative stress, and an abiotic elicitor. *Plant*  
893 *Cell* 10: 359–370.

894 Zhao Y. *et al.* (2015) A metabolomics study delineating geographical location-associated  
895 primary metabolic changes in the leaves of growing tobacco plants by GC-MS and  
896 CE-MS. *Nature* 9: 1–11.

897 **FIGURE LEGENDS**

898 **Figure 1** Total biomass (A), shoot dry weight (B), root dry  
899 weight (C), and root to shoot dry weight ratio (D) on day 29 of  
900 the experiment (n = 10 for all groups). Provenances are:  
901 Rothiemurchus, Scotland (RM), Pernitz, Austria (PER), Jarocin,  
902 Poland (JAR), and Sierra Nevada, Spain (SN). Boxplots  
903 represent the median of the data and the lower and upper  
904 quartiles (25% and 75%). Whiskers represent the most extreme  
905 data point that is no more than 1.5 times the interquartile range  
906 from the box. Control treatment is grey while drought treatment  
907 is black. Different letters indicate significant differences  
908 ( $p < 0.05$ ).

909 **Figure 2** **Figure 2A)** Mean  $\pm$  standard errors for principle  
910 component 1 and **B)** principle component 2 for different  
911 provenances at different time points in the drought experiment  
912 ( $t_1=0$ ,  $t_2=11$  and  $t_3=29$  days). These two components together  
913 explain 20% of the variation in the metabolomics data. Drought  
914 treatment is indicated by the dashed lines. Provenances are  
915 colour coded as follows: blue – Rothiemurchus; orange –  
916 Pernitz; green – Jarocin; red – Sierra Nevada. **C)** Schematic  
917 showing the number of metabolites (out of 4640) that differ  
918 significantly in abundance between given time points and  
919 treatments. The thickness of the arrows between ellipses reflects  
920 the level of the differences in metabolic composition between  
921 treatments and time points.

922 **Figure 3** Volcano plot of the metabolite changes at time point 3  
923 (day 29) after imposing a severe drought. Metabolites are plotted  
924 as a function of the  $\log_2$ (fold change) from control and as –  
925  $\log_{10}$ (P value) of the change from control. Thus, positive values  
926 for fold change indicate increased abundance under drought and  
927 negative values indicated decreased abundance. Metabolites that  
928 showed a  $\log_2$ (fold change) smaller than  $\pm 1$  and a q-value  
929 smaller than 0.05 were selected for further scrutiny and

930 identification (colored in red). Non-selected metabolites are  
931 coloured grey. Note that the volcano plot displays the raw,  
932 uncorrected, p-values. The horizontal dashed line shows the  
933 FDR-corrected critical p-value ( $q < 0,05$ ), i.e. all points with p-  
934 values smaller than the critical value (all points above the line)  
935 are the features for which the null hypothesis of no difference  
936 was rejected. The plot is partially annotated for the compounds  
937 showing the most pronounced and most significant changes  
938 immediately prior to the drought-induced mortality. Legend:  
939 Trp, tryptophan; Tyr, tyrosine; Umb, umbelliferone; Gluc,  
940 dihydrozeatin riboside glucoside; Pro, proline; Phe,  
941 phenylalanine; Skat, skatole; Ser, serine; Jasm, jasmonate; Leu,  
942 leucine; Leu-Gly-Pro, leucine, glycine, proline; Gib,  
943 gibberelline; Styr, styrene; Pyr, pyrole; Spong,  
944 spongipregnoside; Orien, iso-orientin; Ecdys,  
945 dehydroecdysone; Thap, thapsigargin; Osc, oscillatoxin; Spin,  
946 gentibioside spinosin; Cort, hydrocortisone caproate; Purp,  
947 purpurestin; Vit D, vitamine D; Acr, Acronycine.

948

949

950

951

952